

THE TIME COURSE OF ACID-BASE BALANCE  
WHILE ON FBM PATROL

by

Alexander S. Peck, Jr.  
Lieutenant, MC, U.S. Naval Reserve

NAVAL SUBMARINE MEDICAL RESEARCH LABORATORY  
NAVAL SUBMARINE MEDICAL CENTER  
REPORT NUMBER 675

Bureau of Medicine and Surgery, Navy Dept.,  
Research Work Unit MR011.01-5033.01

Reviewed and Approved by:




Charles F. Gell, M.D., D.Sc (Med)  
Scientific Director  
NavSubMedResLab

Reviewed and Approved by:



Joseph D. Bloom, CDR MC USN  
Officer-in-Charge  
NavSubMedResLab

Approved and Released by:



J. E. Stark, CAPT MC USN  
Commanding Officer  
Naval Submarine Medical Center

Approved for public release; distribution unlimited.

## SUMMARY

### THE PROBLEM

To determine the effects of elevated atmospheric CO<sub>2</sub> on arterialized and venous pH, P<sub>CO<sub>2</sub></sub>, and P<sub>O<sub>2</sub></sub>, and to monitor any changes in plasma electrolytes during chronic operational exposure to CO<sub>2</sub>, during an extended FBM submarine patrol.

### FINDINGS

No evidence of respiratory acidosis or change in electrolyte concentration was demonstrated during the study. Calcium and phosphate excretion were also found to be within normal limits with the subjects on unrestricted normal diets.

### APPLICATIONS

This report is of value to all Submarine Medical Officers and environmental physiologists and all personnel who are concerned with the habitability of closed environments over long time periods.

### ADMINISTRATIVE INFORMATION

This investigation was conducted as a part of project MR011. 01-5033, Time Course of Acid-Base Balance While on FBM Patrol. The present report is No. 1 on this project. The manuscript was approved for publication 28 July 1971, and designated as Submarine Medical Research Laboratory Report No. 675.

PUBLISHED BY NAVAL SUBMARINE MEDICAL CENTER

## ABSTRACT

Measurements of arterialized capillary, and venous pH,  $P_{CO_2}$ , and  $P_{O_2}$  were taken from 15 healthy male subjects during a control period and during days 2, 9, 17, 42 and 56 of a Polaris submarine patrol. Venous plasma concentrations of sodium, chloride, and potassium were determined from frozen samples.

Twenty-four hour urine specimens were collected from four of the men in the study. Daily excretion of sodium, potassium, calcium, and phosphate were determined by analysis of frozen samples. The men were on normal diets with no restrictions or control of calcium intake.

Despite prolonged exposure to elevated levels of  $CO_2$  (.72% to .95% with a mean of .85%), no significant respiratory acidosis was documented. Plasma electrolytes were clinically normal and gave no evidence of acidosis during the study or the post-study recovery period. Urinary excretion of calcium, phosphate, sodium, and potassium were within normal limits.

The direct on-board measurement of blood gas and pH values showed little physiological change at present operational levels of  $CO_2$ .

#59/150.

A-W  
A CO<sub>2</sub>, O<sub>2</sub>  
AMINO  
ACIDS

A-A GRADIENT

urine  
PLASMA }

PACKAGE

BIOLOGICAL

NORMAN MACBETH

DATA: of PA. UNDER

# THE TIME COURSE OF ACID-BASE BALANCE WHILE ON FBM PATROL

## INTRODUCTION

The production of compensated respiratory acidosis in man and animals when breathing elevated levels of  $\text{CO}_2$  is a well known and has been amply demonstrated.<sup>2,3</sup> The studies reported by Schaefer et al, were performed with  $\text{CO}_2$  levels up to 1.5%.<sup>1</sup> These studies point out the need for accurate determination of acid-base changes while breathing lower levels of  $\text{CO}_2$ . Additionally, previous determinations have used data derived from frozen samples, rather than measurements of pH,  $\text{PCO}_2$ , and  $\text{PO}_2$ , made at the time of blood collection.<sup>4,5</sup>

This study was designed to document actual, rather than calculated changes in pH,  $\text{PCO}_2$ , and  $\text{PO}_2$  from crew members of an FBM submarine during an operational patrol wherein the  $\text{CO}_2$  level normally averages less than 1%. The electrolyte determinations were carried out on frozen samples of the same blood to serve as confirmation of acidosis, if found by the direct methods. Also of interest were the changes in urinary calcium and inorganic phosphate excretion demonstrated in acidosis. Therefore, it was planned to monitor calcium and phosphate excretion.<sup>6</sup>

## METHODS

Subjects were fifteen healthy volunteer members of an SSBN submarine crew. Fourteen men were between the ages of 23 and 33 years, and one man

was 47 years old. All were in good health, and had no history of pulmonary, renal, or skeletal disease. One man, however, was 11 years post-nephrectomy for correction of a congenital defect. All subjects were exposed to the same operational submarine atmosphere. Oxygen, carbon dioxide and carbon monoxide were measured at four hour intervals in three ship compartments and average values were recorded. On a daily basis, the level of  $\text{CO}_2$  varied from .72% to .95%, with a mean of .85%. Oxygen concentration was maintained between 19% and 21%. Carbon monoxide was less than 25 p.p.m. No dietary restrictions or measurements were in effect during the study. Drinking water was distilled and no minerals were added. At no time during the data collection was the submarine environment ventilated with outside air.

Samples were collected before submerging and during days 2, 9, 17, 30, 42 and 56 of patrol. Recovery samples were obtained 43 days after returning from patrol. Arterialized capillary samples were obtained by digital puncture of siliconized skin after a hand soak of five minutes in water at a temperature of 55°C. All venous samples were collected in heparinized glass vacuum tubes. After measurements of pH,  $\text{PCO}_2$  and  $\text{PO}_2$  were completed, venous samples were immediately centrifuged at 4000 RPM. The plasma was stored at -15°C in capped syringes. The measurements of pH,  $\text{PCO}_2$  and  $\text{PO}_2$  were made with an Instrumentation Laboratory Model 113-SL Ultra Micro

pH Blood Gas Analyzer. Blood bicarbonate values were calculated using a standard Nomogram. Plasma sodium, potassium, and chloride, as well as urinary sodium, potassium, calcium, and phosphate were determined using standard clinical laboratory methods. The recovery values of the pH,  $P_{CO_2}$  and  $P_{O_2}$  were measured on a different instrument of the same model as the instrument used in the on-board measurements. All measurements were corrected for temperature and atmospheric pressure variations.

The urinary excretion of sodium, potassium, and phosphate was monitored by collecting 24-hour urine specimens from four of the men. No control data was collected due to operational limitations. Recovery data was collected and might have been influenced by post patrol dietary changes.

## RESULTS

Table I summarizes the changes in acid-base balance measured during the study. Statistical analysis consisted of a computerized T-test. No definite pattern of respiratory acidosis was demonstrated. The pH values remained essentially constant during the study, while recovery values were elevated significantly when compared with control values. The  $P_{CO_2}$  values showed a similar pattern with a significant depression in arterialized values during day 17. During the study, all pH and  $P_{CO_2}$  values remained within normal limits for healthy males. Blood bicarbonate levels showed significant de-

pression on day 17 in both arterialized and venous sample values. Venous bicarbonate values were significantly depressed on days 42 and 56.

Table II demonstrates essentially constant values of plasma electrolyte concentrations. This would be the expected result if no acidosis were found. Plasma water was also within normal limits.

The excretion of calcium and phosphate tabulated in Table III shows only normal dietary variations and all values fall within accepted normal limits. Accurate statistical analysis was not possible because control data was lacking. It must be noted, however, that due to sample storage space limitations, only four of the fifteen men were monitored for urinary electrolytes and calcium phosphate excretion. Urinary sodium and potassium also were within normal limits.

## DISCUSSION

During the period of this study, current operational levels of  $CO_2$  failed to produce a pattern of respiratory acidosis. Instrumentation was reliable, with only an occasional  $CO_2$  membrane failure. This may account for a reduction in the  $P_{CO_2}$  and bicarbonate data available from several days, especially day 17. The statistical significance determined for day 17 is therefore suspect due to low N values.

In general, venous samples were technically easier to handle and analyze than the arterialized samples and gave more consistent results. The collection

Table I. Arterialized and Venous Blood Levels of pH, P<sub>CO</sub><sub>2</sub>, P<sub>O</sub><sub>2</sub>, (HC03)<sup>-</sup>

	Control	Day 2	Day 9	Day 17	Day 30	Day 42	Day 56	Rec
(Art) pH	X Sem N 7.459 .010 15	7.433 .007 14	7.450 .006 15	7.458 .010 13	7.457 .005 15	7.442 .006 14	7.421 .007 14	7.429 .006 15
(Ven) pH	X Sem N 7.415 .011 15	7.415 .006 14	7.403 .010 15	7.409 .014 13	7.421 .008 15	7.418 .006 14	7.390 .003 14	7.381 .007 15
(Art) P <sub>CO</sub> <sub>2</sub> mm/Hg	X Sem N 36.8 .8 14	37.4 1.0 14	37.6 1.4 10	32.1 1.7 7	37.3 1.2 9	34.8 1.0 8	37.6 1.4 8	41.3 .7 15
(Ven) P <sub>CO</sub> <sub>2</sub> mm/Hg	X Sem N 46.4 1.3 13	46.8 .9 13	46.6 1.6 14	42.9 1.3 10	44.2 1.2 9	42.0 1.6 11	44.3 1.2 8	57.9 1.6 15
(Art) P <sub>O</sub> <sub>2</sub> mm/Hg	X Sem N 87.6 2.0 15	85.6 1.8 14	86.2 2.5 15	97.7 2.6 13	90.8 2.2 15	103.9 4.3 14	103.3 3.1 14	86.1 1.9 15
(Ven) P <sub>O</sub> <sub>2</sub> mm/Hg	X Sem N 35.9 2.7 14	36.9 2.1 14	30.1 2.3 15	33.3 3.8 13	33.7 2.4 15	40.4 3.5 13	42.0 4.0 13	40.0 1.9 15
(Art) (HC03) <sup>-</sup> meq/L	X Sem N 25.8 .7 14	24.8 .7 14	25.8 .8 10	21.7 .9 7	25.6 1.2 10	23.5 .9 8	24.6 .7 8	26.9 .5 15
(Ven) (HC03) <sup>-</sup> meq/L	X Sem N 29.8 .9 13	30.0 .7 13	28.8 1.1 12	26.5 .8 10	28.6 1.0 10	26.9 .9 10	26.5 .6 8	30.3 .7 15

\*Statistically significant at the 5% level.  
Sem -- Standard Error of the mean.

Table II. Plasma Electrolytes and H<sub>2</sub>O (Mean Values) N = 15

	Control	Day 2	Day 9	Day 17	Day 30	Day 42	Day 56	Rec	Expected Normal
Sodium Meq/L	139.3	141.3	138.7	139.0	137.4	139.1	138.6	139.1	136 - 145
Chloride Meq/L	101.1	101.5	101.9	101.6	101.3	101.4	101.2	99.8	100 - 106
Potassium Meq/L	3.9	3.9	3.7	3.4	3.8	3.8	3.7	4.4	3.5 - 5.0
H <sub>2</sub> O%	91.3	90.6	91.2	91.5	91.2	91.0	91.1	91.1	90 - 92

Table III. Twenty-Four Hour Urine Excretion

	Day 2	Day 9	Day 17	Day 30	Day 42	Day 56	Rec	Expected Normal
Sodium Meq/24 hr.	166.5 10.2 4	183.7 10.9 3	174.5 33.3 4	184.0 55.5 3	156.5 25.5 4	234.5 63.2 4	218.0 75.3 4	130 - 260
Potassium Meq/24 hr.	67.3 16.4 4	81.8 26.8 3	78.9 12.8 4	97.9 21.6 3	69.4 15.3 4	82.8 26.4 4	103.9 59.3 4	25 - 100
Calcium Meq/24 hr.	285.0 60.0 4	146.5 51.7 3	166.0 51.0 4	248.0 82.6 3	151.0 43.7 4	193.3 67.8 4	199.0 13.1 4	141 - 364
Phosphate Mg/24 hr. (inorganic)	1426.0 479.0 4	1537.0 576.0 3	1247.0 486.0 4	1457.0 638.0 3	1595.0 785.0 4	1784.0 747.0 4	2590.0 981.0 4	900 - 1300

Ca / mg / 24 hrs

184

184

MEG / L



of blood from fingertip puncture occasionally resulted in air contamination of capillary tubes during measurement. Using ultramicro equipment, air may cause oxygen levels to be abnormally high, and may explain why arterialized samples had slightly higher oxygen tensions on several days. The use of ear lobe puncture for arterialized samples may be adviseable in future work.

All recovery samples were performed on a different instrument and by a different technician than were the on-board values. This may have introduced some small constant error in pH,  $PCO_2$  and  $PO_2$  data. Slight changes in calibration and individual technique are to be expected.

Plasma electrolyte concentrations remained constant and no chloride changes were seen that were compatible with acidosis. Since only mild changes were demonstrated at 1.5%  $CO_2$  levels, submarines could be operating with a concentration of  $CO_2$  which fails to produce respiratory acidosis. There are several other factors which must be investigated under low levels of atmospheric  $CO_2$ . These factors include: (a) individual response to various concentrations of carbon dioxide, and (b) the effect of circadian biological cycles on the overall acid-base changes.

In the past, several attempts have been made to classify men into categories of "responders" and "non-responders" when exposed to elevated levels of  $CO_2$ .<sup>7</sup> An initial attempt at separating our volunteers into groups according to  $CO_2$  tolerance has been started at this Laboratory. At the present time, a method for definitive

evaluation of each individual is not yet available.

It is well known that cyclic biological functions can cause changes in calcium metabolism and may affect acid-base balance as well. Blood samples were drawn during the work cycle of the individual volunteer. It was not possible to collect samples on a fixed, pre-determined time schedule due to rotation of watches and other operational commitments.

It is concluded that the results of this study would seem to substantiate the adequacy of present standards of atmosphere control. No significant physiological changes were demonstrated under the operational levels of  $CO_2$  maintained on this particular submarine. Calcium and phosphate excretions were determined to be within normal limits with variation due to diet. Accurate dietary control is possible, but difficult to enforce on an operational submarine.

#### REFERENCES

1. Schaefer, K. E., Nichols, G. and Carey, C. R.: Acid-base balance and blood and urine electrolytes of man during acclimatization to  $CO_2$ . J. Appl Physiol, Vol. 19, No. 1, January 1964.
2. Schaefer, K. E.: Respiration and acid-base balance during prolonged exposure to 3  $CO_2$ . Pflueg. Arch. Ges. Phys., 251:689.
3. Schaefer, K. E., Messier, A. A. and Morgan, C.: Effect of inter-

mittent exposure to 3% CO<sub>2</sub> on acid-base balance. Naval Submarine Medical Research Laboratory, Report No. 635, Naval Submarine Medical Center, Groton, Conn.

4. Gortner, D. A.: Effects of elevated atmospheric CO<sub>2</sub> on acid-base balance and plasma and red cell electrolytes on FBM submarine crew members. Thesis report, Naval Submarine Medical Center, Groton, Connecticut.
5. Gray, S. P., Lambert, R. J. W., and Morris, J. E.: Calcium magnesium and phosphate metabolism during prolonged exposure to carbon dioxide. 15th Meeting R.N. Med. Aerospace Council, Submarine Sub-Committee, England 1968.
6. Schaefer, K. E., Nichols, G. and Carey, C. R.: Calcium phosphorus metabolism in man during acclimatization to carbon dioxide. J. Appl. Physiol., Vol. 18, No. 6, November 1963.
7. Schaefer, K. E., and Dougherty, J. H.: Variability of Respiratory Functions Based on Circadian Cycles. National Aeronautics and Space Administration and Naval Submarine Medical Center Report No. 486, November, 1966.

UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R & D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1. ORIGINATING ACTIVITY (Corporate author) NAVAL SUBMARINE MEDICAL CENTER, Submarine Medical Research Laboratory		2a. REPORT SECURITY CLASSIFICATION Unclassified
		2b. GROUP
3. REPORT TITLE  THE TIME COURSE OF ACID-BASE BALANCE WHILE ON FBM PATROL.		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Report on in-house project.		
5. AUTHOR(S) (First name, middle initial, last name)  Alexander S. PECK, JR., LT, MC, USNR		
6. REPORT DATE 27 July 1971	7a. TOTAL NO. OF PAGES 6	7b. NO. OF REFS 7
8a. CONTRACT OR GRANT NO.	9a. ORIGINATOR'S REPORT NUMBER(S) Naval Submarine Medical Research Lab. Report No. 675	
b. PROJECT NO. MR011.01-5033.01		
c.	9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d.		
10. DISTRIBUTION STATEMENT  Approved for public release; distribution unlimited.		
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY Naval Submarine Medical Center Box 600 Naval Submarine Base Groton, Connecticut 06340
13. ABSTRACT  Measurements of arterialized capillary, and venous pH, $PCO_2$ and $P_{O_2}$ were taken from 15 healthy male subjects during a control period and during days 2, 9, 17, 42 and 56 of a Polaris submarine patrol. Venous plasma concentrations of sodium, chloride, and potassium were determined from frozen samples. Twenty-four hour urine specimens were collected from four of the men in the study. Daily excretion of sodium, potassium, calcium, and phosphate were determined by analysis of frozen samples. The men were on normal diets with no restrictions or control of calcium intake. Despite prolonged exposure to elevated levels of $CO_2$ (.72% to .95% with a mean of .85%), no significant respiratory acidosis was documented. Plasma electrolytes were clinically normal and gave no evidence of acidosis during the study and post recovery period. Urinary excretion of calcium, phosphate, sodium, and potassium were within normal limits. The direct on-board measurement of blood gas and pH values shows little physiological change at present operational levels of $CO_2$ .		

DD FORM 1 NOV 65 1473 (PAGE 1)

S/N 0102-014-6600

UNCLASSIFIED

Security Classification

UNCLASSIFIED

Security Classification

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Carbon dioxide exposures. Polaris. Plasma electrolytes. Respiratory acidosis. Calcium excretion. Phosphate excretion.						

UNCLASSIFIED

Security Classification